

## Fock Space: Second quantization representation

Let us define the **creation operators** acting on the vectors in the Fock space:

$$a_{\mu}^{+} \Phi_{\mu_1 \dots \mu_A} \equiv \begin{cases} 0 & \text{if } \mu \in \{\mu_i\} \\ \Phi_{\mu \mu_1 \dots \mu_A} & \text{if } \mu \notin \{\mu_i\} \end{cases}$$

$a_{\mu}^{+}$  takes us from  $\mathcal{H}_A$  to  $\mathcal{H}_{A+1}$

By calculating the scalar product of  $a_{\mu}^{+} \Phi_{\nu_1 \dots \nu_A}$

with an arbitrary state  $\Phi_{\mu_1 \dots \mu_{A+1}}$  it is easy to show that, for

$$a_{\mu} \equiv (a_{\mu}^{+})^{+}$$

$$a_{\mu} \Phi_{\mu_1 \dots \mu_{A+1}} = \begin{cases} 0 & \text{if } \mu \notin \{\mu_i\} \\ (-1)^{k+1} \Phi_{\mu_1 \dots \mu_{k-1} \mu_{k+1} \dots \mu_{A+1}} & \text{if } \mu = \mu_k \end{cases}$$

↑  
 $\mu$  is absent!

$a_{\mu}$  takes us from  $\mathcal{H}_{A+1}$  to  $\mathcal{H}_A$

... thus is called an “**annihilation operator**”

let us consider  $a_\mu^+$  and  $a_\nu^+$  ( $\mu \neq \nu$ )

and  $\Phi_{\mu_1 \dots \mu_A}$  ( $\mu \neq \mu_i, \nu \neq \mu_i$ )

$$\begin{aligned} a_\mu^+ a_\nu^+ \Phi_{\mu_1 \dots \mu_A} &= \Phi_{\mu\nu\mu_1 \dots \mu_A} = \\ &= -\Phi_{\nu\mu\mu_1 \dots \mu_A} = -a_\nu^+ a_\mu^+ \Phi_{\mu_1 \dots \mu_A} \end{aligned}$$

$$\begin{aligned} \longrightarrow \quad a_\mu^+ a_\nu^+ &= -a_\nu^+ a_\mu^+ \\ a_\nu a_\mu &= -a_\mu a_\nu \end{aligned}$$

Similarly, for  $\nu=\mu_k$  and  $\nu \neq \mu_j$ , one obtains

$$\begin{aligned} a_\mu^+ a_\nu \Phi_{\mu_1 \dots \mu_A} &= (-1)^{k+1} \Phi_{\mu\mu_1 \dots \mu_{k-1} \mu_{k+1} \dots \mu_A} = \\ &= -a_\nu \Phi_{\mu\mu_1 \dots \mu_A} = -a_\nu a_\mu^+ \Phi_{\mu_1 \dots \mu_A} \end{aligned}$$

For  $\nu=\mu=\mu_k$ :

$$\begin{aligned} a_\nu^+ a_\nu \Phi_{\mu_1 \dots \mu_A} &= (-1)^{k+1} \Phi_{\nu\mu_1 \dots \mu_{k-1} \mu_{k+1} \dots \mu_A} = \\ &= \Phi_{\mu_1 \dots \mu_A} = \Phi_{\mu_1 \dots \mu_A} - a_\nu a_\nu^+ \Phi_{\mu_1 \dots \mu_A} \end{aligned}$$

$$\longrightarrow \quad a_\mu^+ a_\nu = \delta_{\mu\nu} - a_\nu a_\mu^+$$

$$\begin{aligned}
 \{a_{\mu}^{+}, a_{\nu}^{+}\} &= 0 \\
 \{a_{\mu}, a_{\nu}\} &= 0 \\
 \{a_{\mu}^{+}, a_{\nu}\} &= \delta_{\mu\nu}
 \end{aligned}
 \tag{1}$$

Consequently,  $(a_{\mu}^{+})^2 = 0$ ,  $(a_{\mu})^2 = 0$

**(The Pauli Principle!)**

In addition, we have to consider a case of the vacuum vector:

$$a_{\mu}^{+} \Phi_{vac} = \Phi_{\mu} = \phi_{\mu}(x)$$

$$a_{\mu} \Phi_{vac} = 0$$

The Fock space defines  $a^{+}$  and  $a$  !!!

But...

...the Fock space can be defined through  $a^{+}$  and  $a$  !!!

Namely, given a set of  $M$  operators  $a_{\mu}^{+}$  and  $a_{\mu}$  that meet

the anticommutation rules (1) in a given Hilbert space, and

assuming that there exists a vacuum state  $|0\rangle$ , such that

$$a_{\mu}|0\rangle = 0 \quad \text{for any } \mu$$

... one can show that the set of states

$$|\mu_1 \cdots \mu_A\rangle \equiv a_{\mu_1}^+ \cdots a_{\mu_A}^+ |0\rangle, \quad \mu_1 < \cdots < \mu_A$$

where  $A=0,1,\dots,M$ , is a basis of a space that is isomorphic with the Fock space, that is:

$$\begin{aligned} \Phi_{\mu_1 \cdots \mu_A} &\leftrightarrow |\mu_1 \cdots \mu_A\rangle \\ \Phi_{vac} &\leftrightarrow |0\rangle \end{aligned}$$

The vectors  $|\mu_1 \cdots \mu_A\rangle$  form a basis in the Fock Space (or, more precisely, a basis of the representation of the Fock Space), which is called the **second quantization representation**.

## Properties of the vacuum

1. For a given set of creation and annihilation operators acting in a Fock space, there exists only one vacuum.

Proof: let us assume that in addition to  $|0\rangle$ ,  $\langle 0|0\rangle = 1$ ,

there exists another vacuum  $|0'\rangle$ ,  $\langle 0'|0'\rangle = 1$ ,

such that  $a_\mu|0'\rangle = 0$  for any  $\mu$ . Of course,

$$\langle 0'|a_{\mu_1}^+ a_{\mu_2}^+ \cdots a_{\mu_A}^+ |0\rangle = 0 \quad (A = 1, 2, \cdots M)$$

If  $\langle 0'|0\rangle = 0$  then the vector  $|0'\rangle$  is orthogonal to all the vectors in the Fock space; hence, it has to be a zero vector.

Consequently,  $\langle 0'|0\rangle \neq 0$ .

Let us, therefore consider the vector  $|0'\rangle - \langle 0'|0\rangle|0\rangle$

that is orthogonal to  $|0\rangle$  and is also a vacuum. Hence, it has to be a zero vector. Thus, finally,

$$|0'\rangle = \langle 0'|0\rangle|0\rangle$$

a phase...